hep-th/9912033 UCLA/99/TEP/55

On the Coupling of Gravitons to Matter

Z. Bern, A. De Freitas and H.L. Wong
Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095-1547
(December, 1999)

Using relationships between open and closed strings, we present a construction of tree-level scattering amplitudes for gravitons minimally coupled to matter in terms of gauge theory partial amplitudes. In particular, we present examples of amplitudes with gravitons coupled to vectors or to a single fermion pair. We also present two examples with massive graviton exchange, as would arise in the presence of large compact dimensions. The gauge charges are represented by flavors of dynamical scalars or fermions. This also leads to an unconventional decomposition of color and kinematics in gauge theories.

PACS numbers: 11.15.Bt, 11.25.-w, 11.55.-m, 11.25.Db, 95.30.Sf

arXiv:hep-th/9912033v1 4 Dec 1999

Although gravity and gauge theories both contain local symmetries, their dynamical behavior are rather different. Nonetheless, there are some rather striking relationships between the theories such as the AdS/CFT correspondence [1] and the perturbative relationships implied by the structure of closed string amplitudes in terms of open string amplitudes [2]. In this letter we demonstrate that at least for the case of a flat background, there is an elegant and powerful way to obtain the interaction of gravitons with matter, based on the structure of heterotic strings. Our construction is valid for the cases of gravitons coupled to vectors and to at most a single fermion pair. The cases of pure Einstein gravity and $N = 8$ supergravity have already been discussed in refs. [3,4,5]. Recently, there has been a resurgence of interest in perturbative gravity, especially in the presence of large compact dimensions [6] and in AdS spaces [1], making it all the more timely to investigate the perturbative structure of gravity theories.

The differences between gravitational and gauge interactions are well known. For example, quantum chromodynamics is renormalizable and exhibits confinement. Neither property is shared by gravity. Moreover, the structure of the Einstein-Hilbert and Yang-Mills Lagrangians $\mathcal{L}_{\rm EH} = \frac{2}{\kappa^2}\sqrt{-g}R$ and $\mathcal{L}_{\rm YM} = -\frac{1}{4}F^a_{\mu\nu}F^{\mu\nu a}$ are rather different. In particular, gravity has an infinite set of contact interactions while non-abelian gauge theories have only up to four-point contact interactions.

At tree level, Kawai, Lewellen and Tye (KLT) [2] have, however, expressed closed string amplitudes as sums of products of open string amplitudes. In the field theory limit this allows one to obtain gravity tree amplitudes directly from gauge theory ones. This is generally simpler than using conventional gravity Feynman rules. As discussed in ref. [7], by using non-linear field redefinitions and gauge fixings, the Lagrangians of the two theories can be rearranged to make the relations between the two theories more apparent. (See also ref. [8].) With the analytic reconstruction method described in refs. [9,4,5] the loop amplitudes can be obtained from the tree ones. In this way gravity can be perturbatively quantized without reference to a Lagrangian. This has already been exploited to demonstrate that $D = 11$ supergravity diverges at two-loops in accordance with naive expectations [4,10] and that $D = 4$ $N = 8$ supergravity likely has the first divergence at five [4] instead of three loops [11], as expected from superspace power counting. In order to perform a more general investigation of the divergence structure of gravity theories, it would be useful to have more general tree amplitudes besides those of pure gravity or supergravity.

Here we discuss amplitudes where gravitons are minimally coupled to gauge theories. Maximally helicity violating (MHV) n-point tree amplitudes for pure gravity [3] and for gravitationally dressed Parke-Taylor [12] gluon amplitudes have been presented in refs. [3,13]. Using the KLT relations, we extend these results to the remaining MHV gravitationally dressed gluon amplitudes and to ones with a single fermion pair. Our construction also applies to cases of non-maximal helicity violation, but the MHV cases are the simplest to present. We also present four-point examples with a (massive) graviton exchange.

The KLT equations [2,3] for field theory gravity tree amplitudes are,

$$\begin{aligned}
&\mathcal{M}_4(1,2,3,4) = -is_{12}A_4(1,2,3,4)\tilde{A}_4(1,2,4,3)\,,\\
&\mathcal{M}_5(1,2,3,4,5) =\\
&\quad i\Big[s_{12}s_{34}A_5(1,2,3,4,5)\tilde{A}_5(2,1,4,3,5)\\
&\qquad +s_{13}s_{24}A_5(1,3,2,4,5)\,\tilde{A}_5(3,1,4,2,5)\Big]\,,
\end{aligned} \tag{1}$$

where $s_{ij} = (k_i + k_j)^2$. The A_n and $\tilde{A}_n$ are tree-level color-ordered gauge theory partial amplitudes [14]. Our notation is to suppress polarizations, ε_i, and momenta, k_i, leaving only the label i for each leg. Explicit forms of the KLT equations for an arbitrary number of external

legs may be found in ref. [5]. We have dropped the couplings, but for each gauge and gravitational interaction appearing on the left-hand-side, factors of the couplings g and $\kappa/2$ (where $\kappa^2 = 32\pi G_N$) are respectively required.

Although the use of the KLT equations in the field theory limit does not require a fully consistent string theory, some features of string theory are retained. For example, an oriented string will necessarily contain antisymmetric tensor and dilaton states in its massless spectrum. When describing graviton exchanges, the KLT equations will therefore also implicitly contain these states. However, for the cases dealt with in this letter, the antisymmetric tensor and dilaton decouple, so that extra projections are not required to remove these states, if unwanted.

In order to apply the KLT equations to the cases of gravity coupled to matter, we must specify the factorization of each state into states appearing solely in gauge theories. In doing so, we must ensure that the spin and charges for each state are the desired ones. We factorize each graviton into a product of vectors, each gluon into either a product of a vector and a scalar or a product of two fermions and each quark into a fermion and a scalar. As motivated by the structure of the heterotic string, the gauge charges of each physical state are assigned as flavor charges carried by scalars or by fermions. At the linearized level these factorizations are trivial. In a sense, we use the KLT equations to extend them to the fully interacting theories.

As a specific example, consider $SU(N_c)$ gauge theory minimally coupled to Einstein gravity. In fig. 1 we present rules for constructing, via the KLT equations, gluon amplitudes dressed by minimally coupled gravitons; for this case, the rules give the partial amplitudes A_n and $\tilde{A}_n$ appearing in the KLT equations. These rules are determined by requiring that the amplitudes with gravitons carry the correct quantum numbers and dimensions. As discussed below, cases with a single fermion pair can be handled similarly. (The signature of $\eta_{\mu\nu}$ is $(+,-,-,-)$.) These rules are similar to ordinary gauge theory Feynman rules except that they are color ordered [14], i.e. the color group theory factors have been stripped. In constructing the diagrams one should maintain a clockwise ordering of the legs. The group theory factors appearing in the rule are associated with the flavors of dynamical scalars. After substituting the A_n and $\tilde{A}_n$ into the KLT equations, on the left-hand-side the group theory factors are reinterpreted as color factors. We have displayed the vertices in the commonly used Feynman gauge and have chosen all group theory factors to be in the adjoint representation. For other representations the group theory factors should be altered to the appropriate one.

It is instructive to first apply these rules to a pure gluon amplitude. The KLT equations will apply, since such amplitudes are part of a theory coupling gravitons to gluons. In this case, the construction is circular, since



FIG. 1. The color-ordered Feynman vertices for obtaining, via the KLT equations, gluon amplitudes dressed with gravitons. Dashed lines represent scalars and curly ones vectors.

it starts with a gauge theory amplitude only to wind up at the same amplitude. It will, however, easily generalize to cases including gravitons.

With the standard color decomposition [14], the four-gluon tree amplitude is expressed as,

$$\mathcal{M}_4(1_g^{a_1}, 2_g^{a_2}, 3_g^{a_3}, 4_g^{a_4}) = g^2 \sum_\sigma \mathrm{Tr}[T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} T^{a_{\sigma(3)}} T^{a_{\sigma(4)}}] \times A_4(\sigma(1_g), \sigma(2_g), \sigma(3_g), \sigma(4_g))\,, \qquad (2)$$

where σ runs over the set of non-cyclic permutations and the A_4 are color-ordered partial amplitudes. We denote gluon, graviton, scalar and quark legs by the subscripts g, h, s and q. (Our fundamental representation color matrices are normalized by $\mathrm{Tr}[T^a T^b] = \delta^{ab}$.)

By applying the KLT formula (1) we express the four-gluon amplitude with only gluon exchanges as

$$\mathcal{M}_4(1_g^{a_1}, 2_g^{a_2}, 3_g^{a_3}, 4_g^{a_4}) = -ig^2 s_{12} A_4(1_g, 2_g, 3_g, 4_g) \times \tilde{A}_4(1_s^{a_1}, 2_s^{a_2}, 4_s^{a_4}, 3_s^{a_3})\,, \qquad (3)$$

where the diagrams for obtaining $A_4(1_g, 2_g, 3_g, 4_g)$ and $\tilde{A}_4(1_s, 2_s, 4_s, 3_s)$ are given in figs. 2 and 3. Note that in each of the two cases the labels on the diagrams follow the cyclic ordering of the arguments of the amplitudes. The value of the first of these partial amplitudes is well known and may be found in, for example, ref. [14]. Evaluating the diagrams in fig. 3, the four-scalar partial amplitude is,

$$\tilde{A}_4(1_s^{a_1}, 2_s^{a_2}, 4_s^{a_4}, 3_s^{a_3}) = 2i\frac{f^{a_1 a_2 b} f^{a_4 a_3 b}}{s_{12}} + 2i\frac{f^{a_2 a_4 b} f^{a_3 a_1 b}}{s_{13}}\,,$$

where $f^{abc} = -i\,\mathrm{Tr}\big([T^a, T^b]T^c\big)/\sqrt{2}$ are the group structure constants. This type of decomposition of group theory and kinematics is rather unconventional. Nevertheless, it is straightforward to verify that the amplitude in eq. (3) is equal to that in eq. (2). Similar decompositions hold for any number of external legs, by applying

n-point KLT equations. In this way both graviton and full gauge theory amplitudes satisfy KLT equations, providing a natural way to obtain mixed amplitudes of states appearing in gravity and in gauge theories.



FIG. 2. The color-ordered Feynman diagrams contributing to $A(1_g, 2_g, 3_g, 4_g)$.



FIG. 3. The two diagrams contributing to the scalar partial amplitude $\tilde{A}_4(1_s^{a_1}, 2_s^{a_2}, 4_s^{a_4}, 3_s^{a_3})$. Each diagram is dressed with flavor group theory factors.

For example, we can express $\mathcal{M}_4(1_g, 2_g, 3_g, 4_h)$, in terms of the gauge theory partial amplitudes $A_4(1_g, 2_g, 3_g, 4_g)$, whose diagrams are given in fig. 2, and the three-scalar one-vector partial amplitude $\tilde{A}_4(1_s^{a_1}, 2_s^{a_2}, 4_g, 3_s^{a_3})$ whose diagrams are the same as those in fig. 3, except that leg 4 is replaced with a vector.

More generally, by applying the Feynman rules in fig. 1 we have obtained four-, five-, and six-point mixed amplitudes for gluons and gravitons. A sampling of these is,

$$M_5(1_g^+, 2_g^+, 3_g^+, 4_g^-, 5_h^-) = -ig^2\Big(\frac{\kappa}{2}\Big)\frac{\langle 4\,5\rangle^4}{\langle 1\,2\rangle\langle 2\,3\rangle\langle 3\,4\rangle\langle 4\,1\rangle}\,,$$

$$M_5(1_g^-, 2_g^+, 3_g^+, 4_h^+, 5_h^-) = ig\Big(\frac{\kappa}{2}\Big)^2\frac{\langle 1\,5\rangle^4}{\langle 1\,2\rangle\langle 2\,3\rangle\langle 3\,1\rangle} \times\frac{1}{\langle 4\,1\rangle\langle 4\,5\rangle}\Big[\langle 2\,1\rangle\langle 2\,5\rangle\frac{[2\,4]}{\langle 2\,4\rangle} + \langle 3\,1\rangle\langle 3\,5\rangle\frac{[3\,4]}{\langle 3\,4\rangle}\Big]\,,$$

$$M_5(1_g^+, 2_g^+, 3_h^+, 4_h^-, 5_h^-) = 0\,,$$

$$M_5(1_g^-, 2_g^+, 3_h^+, 4_h^+, 5_h^-) = i\Big(\frac{\kappa}{2}\Big)^3\langle 1\,5\rangle^4\langle 2\,5\rangle^2 \times h(1, \{2,3,4\}, 5)\,,$$

for the coefficient of the color traces $\mathrm{Tr}[T^{a_i}\cdots T^{a_j}]$ which follow the ordering of gluon legs. The complete amplitudes are given by summing over non-cyclic permutations of the gluon legs after multiplying by the color traces. The function $h(a, \{P\}, b)$ appearing in the last amplitude is the half-soft function defined in ref. [5]. Our convention for the '$\pm$' helicity labeling is to take all legs to be outgoing. In these amplitudes graviton exchanges between gluons are not incorporated; such contributions are suppressed by additional powers of the gravitational coupling. We have expressed the amplitudes in terms of $D = 4$ spinor inner products. (See e.g. ref. [14].) These are denoted by $\langle i\,j\rangle = \langle i^-|j^+\rangle$ and $[i\,j] = \langle i^+|j^-\rangle$, where $|i^\pm\rangle$ are massless Weyl spinors of momentum k_i, labeled with the sign of the helicity. They are antisymmetric, with norm $|\langle i\,j\rangle| = |[i\,j]| = \sqrt{s_{ij}}$.

Although $D = 4$ spinor helicity is quite useful, the KLT equations do not require its use, and are valid in arbitrary dimensions. Likewise, the conditions of maximal helicity violation are not essential, although they considerably simplify amplitudes. We have, for example, obtained all the independent amplitudes with five gluons and one graviton using the KLT equations and the six-gluon partial amplitudes given in ref. [14].

The cases corresponding to gravitationally dressed Parke-Taylor amplitudes match those previously computed by Selivanov [13] using a generating function method. Selivanov's n-point expression for the coefficient of $\mathrm{Tr}[T^{a_1}T^{a_2}\cdots T^{a_m}]$ is

$$M_n(1_g^-, 2_g^+, \ldots, i_g^-, \ldots, m_g^+, (m+1)_h^+, \ldots, n_h^+) = ig^{m-2}\Big(-\frac{\kappa}{2}\Big)^{n-m}\frac{\langle 1\,i\rangle^4}{\langle 1\,2\rangle\langle 2\,3\rangle\cdots\langle m\,1\rangle}S(1, i, \{h^+\}, \{g^+\})\,,$$

where all legs have positive helicity except 1 and i and

$$S(i, j, \{h^+\}, \{g^+\}) = \Big(\prod_{m\in\{h^+\}}\frac{d}{da_m}\Big) \times\prod_{l\in\{g^+\}}\exp\Big[\sum_{n_1\in\{h^+\}}a_{n_1}\frac{\langle l\,i\rangle\langle l\,j\rangle[l\,n_1]}{\langle n_1\,i\rangle\langle n_1\,j\rangle\langle l\,n_1\rangle} \times\exp\Big[\sum_{\substack{n_2\in\{h^+\}\\ n_2\neq n_1}}a_{n_2}\frac{\langle n_1\,i\rangle\langle n_1\,j\rangle[n_1\,n_2]}{\langle n_2\,i\rangle\langle n_2\,j\rangle\langle n_1\,n_2\rangle}\exp\Big[\ldots\Big]\Big]\Big]\Big|_{a_j=0}\,. \tag{4}$$

In this factor legs i and j are the negative helicity ones while the sets $\{h^+\}$ and $\{g^+\}$ represent the positive helicity graviton and gluon legs.

The KLT equations immediately show that mixed gluon and graviton amplitudes vanish if all legs have identical helicity or if there is a single leg of opposite helicity, since the corresponding pure gluon [14] amplitudes vanish. We also extrapolate our results to n-legs to obtain the remaining non-vanishing MHV gluon amplitudes dressed with gravitons. Doing so yields,

$$M_n(1_g^-, 2_g^+, 3_g^+, \ldots, m_g^+, (m+1)_h^-, (m+2)_h^+, \ldots, n_h^+) = ig^{m-2}\Big(-\frac{\kappa}{2}\Big)^{n-m}\frac{\langle 1\;m+1\rangle^4}{\langle 1\,2\rangle\langle 2\,3\rangle\cdots\langle m\,1\rangle} \times S(1, m+1, \{h^+\}, \{g^+\})\,, \tag{5}$$

for the coefficient of the color factor $\mathrm{Tr}[T^{a_1}T^{a_2}\cdots T^{a_m}]$. Furthermore, the amplitudes where the gluons are all of identical helicity vanish, independent of the graviton helicity configuration. These amplitudes were constructed by demanding that they satisfy the required factorization properties in all channels: The amplitudes have no multi-particle poles, obey the collinear and soft factorization for gluons given in ref. [14] and the ones for gravitons

given in ref. [5]. Moreover, they have the correct phase singularity for a gluon becoming collinear with a graviton. Although this does not constitute a complete proof that these results are valid for all n, in the past similar analytic requirements have invariably led to the correct results. (See e.g. refs. [12,15].)

The KLT equation (1) can also be directly applied to the case of a single fermion pair, by using the vertex $i\gamma^\mu/\sqrt{2}$ for the coupling of a fermion pair to a vector. For example, in eq. (5) the result of replacing legs 1 and 2 by gluinos is given by multiplying by the overall factor $\langle 2\ m+1\rangle\,/\,\langle 1\ m+1\rangle$. This is consistent with the result obtained by applying $N = 1$ supersymmetry Ward identities [16]. For fundamental representation quarks one must also modify the associated color factor to $(T^{a_3}\cdots T^{a_m})^{\bar{\imath}_2}_{i_1}$. Cases with a single gravitino pair are similar. On the other hand, cases with multiple fermion pairs are more intricate. In particular, for the KLT factorization to work, auxiliary rules for assigning global charges in the color-ordered amplitudes appear to be necessary.

Although, we do not have a general formalism, it is interesting that a KLT factorization can also be found for examples with a massive graviton. For example, the four-gluon amplitude with a massive graviton exchange can be factorized as

$$\begin{aligned}
&\mathcal{M}_4^{\rm ex}(1_g^{-a_1}, 2_g^{+a_2}, 3_g^{-a_3}, 4_g^{+a_4}) \\
&\quad = -i\left(\frac{\kappa}{2}\right)^2 (s_{12}-m^2)\, A_4(1_{\bar q}^{-a_1}, 2_q^{+a_2}, 3_{\bar q}^{-a_3}, 4_q^{+a_4}) \\
&\qquad\qquad \times \tilde{A}_4(1_{\bar q}^-, 2_q^+, 4_q^+, 3_{\bar q}^-) \qquad (6)\\
&\quad = -i\left(\frac{\kappa}{2}\right)^2 [2\,4]^2\,\langle 1\,3\rangle^2 \left(\frac{\delta^{a_1a_2}\delta^{a_3a_4}}{s_{12}-m^2} + \frac{\delta^{a_1a_4}\delta^{a_2a_3}}{s_{14}-m^2}\right).
\end{aligned}$$

On the right-hand-side the color-ordered amplitudes are those for quark scattering via massive vector boson exchanges. (This factorization works because of our particular assignment of a_i charges.) Similarly, the scattering of fundamental representation quarks by gluons via massive graviton exchange is

$$\begin{aligned}
&\mathcal{M}_4^{\rm ex}(1_q^{-i_1}, 2_{\bar q}^{+\bar\imath_2}, 3_g^{-a_3}, 4_g^{+a_4}) \\
&\qquad = -i\Big(\frac{\kappa}{2}\Big)^2 (s_{12}-m^2) A_4(1_s^{i_1}, 2_s^{\bar\imath_2}, 3_Q^{-a_3}, 4_{\bar Q}^{+a_4}) \\
&\qquad\qquad \times \tilde{A}_4(1_q^-, 2_{\bar q}^+, 4_{\bar Q}^+, 3_Q^-) \qquad (7)\\
&\qquad = i\Big(\frac{\kappa}{2}\Big)^2 \langle 1\,3\rangle^2\,[4\,2]\,[4\,1]\,\frac{\delta_{i_1}{}^{\bar\imath_2}\delta^{a_3a_4}}{s_{12}-m^2}\,,
\end{aligned}$$

where q and Q are distinct massless fermion species. For both amplitudes (6) and (7) we have verified that both sides of the equations are equal by explicitly computing the amplitudes on the left-hand-side using conventional Feynman rules.

The examples (6) and (7) suggest that it should be possible to obtain a general formalism for (massive) graviton exchanges. With multiple fermion pairs the antisymmetric tensor and dilaton will decouple only with extra projections.

It would be interesting to see if the relationships between gravity and gauge theories discussed here can be extended to more general field configurations.

We thank Lance Dixon and Maxim Perelstein for useful discussions. This work was supported by the US Department of Energy under grant DE-FG03-91ER40662.

[1] O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, preprint hep-th/9905111.

[2] H. Kawai, D.C. Lewellen and S.-H.H. Tye, Nucl. Phys. B269:1 (1986).

[3] F.A. Berends, W.T. Giele and H. Kuijf, Phys. Lett. 211B:91 (1988).

[4] Z. Bern, L. Dixon, D.C. Dunbar, M. Perelstein and J.S. Rozowsky, Nucl. Phys. B530:401 (1998) [hep-th/9802162].

[5] Z. Bern, L. Dixon, M. Perelstein and J.S. Rozowsky, Phys. Lett. B444:273 (1998) [hep-th/9809160]; Nucl. Phys. B546:423 (1999) [hep-th/9811140].

[6] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B429:263 (1998) [hep-ph/9803315].

[7] Z. Bern and A.K. Grant, Phys. Lett. B457:23 (1999) [hep-th/9904026].

[8] W. Siegel, Phys. Rev. D47:5453 (1993) [hep-th/9302036]; in *Proceedings of Strings 1993*, eds. M.B. Halpern, A. Sevrin and G. Rivlis (World Scientific, Singapore, 1994) [hep-th/9308133].

[9] Z. Bern, L. Dixon and D.A. Kosower, Ann. Rev. Nucl. Part. Sci. 46:109 (1996) [hep-ph/9602280].

[10] S. Deser, D. Seminara, Phys. Rev. Lett. 82:2435 (1999) [hep-th/9812136].

[11] P.S. Howe and K.S. Stelle, Int. J. Mod. Phys. A4:1871 (1989).

[12] S.J. Parke and T.R. Taylor, Phys. Rev. Lett. 56:2459 (1986);

[13] K.G. Selivanov, Phys. Lett. B420:274 (1998) [hep-th/9710197]; Mod. Phys. Lett. A12:3087 (1997) [hep-th/9711111].

[14] M. Mangano and S.J. Parke, Phys. Rep. 200:301 (1991); L. Dixon, in *Proceedings of Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 95)*, ed. D.E. Soper [hep-ph/9601359].

[15] F.A. Berends and W.T. Giele, Nucl. Phys. B306:759 (1988); Z. Bern, G. Chalmers, L. Dixon and D.A. Kosower, Phys. Rev. Lett. 72:2134 (1994), hep-ph/9312333; G.D. Mahlon, Phys. Rev. D49:2197 (1994); Phys. Rev. D49:4438 (1994); Z. Bern, L. Dixon, D.C. Dunbar and D.A. Kosower, Nucl. Phys. B425:217 (1994), hep-ph/9403226.

[16] M.T. Grisaru and H.N. Pendleton, Nucl. Phys. B124:81 (1977); S.J. Parke and T. Taylor, Phys. Lett. 157B:81 (1985).